

Elements of harmonic analysis, 5

Stephen William Semmes
Rice University
Houston, Texas

Fix a positive integer n , and let $\mathcal{S}(\mathbf{R}^n)$ denote the vector space of complex-valued smooth functions $f(x)$ on $\mathbf{R}^n$ such that f and all of its derivatives are bounded functions on $\mathbf{R}^n$, and moreover they are rapidly decreasing in the sense that

$$(1) \quad x^\alpha \frac{\partial^{d(\beta)}}{\partial x^\beta} f(x)$$

are also bounded functions on $\mathbf{R}^n$ for all multi-indices α, β , where $d(\beta)$ denotes the degree of β , which is to say the sum of its components. Thus $\mathcal{S}(\mathbf{R}^n)$ contains the smooth functions on $\mathbf{R}^n$ with compact support as a linear subspace. Also, $\mathcal{S}(\mathbf{R}^n)$ contains the Gaussian functions $\exp(-A(x) \cdot x)$, where A is a positive-definite symmetric linear transformation on $\mathbf{R}^n$, and $x \cdot y$ is the usual inner product defined by

$$(2) \quad x \cdot y = \sum_{j=1}^n x_j y_j.$$

The space $\mathcal{S}(\mathbf{R}^n)$ is commonly known as the *Schwartz class* of rapidly decreasing smooth functions on $\mathbf{R}^n$. If f_1, f_2 are functions in the Schwartz class, then the sum $f_1 + f_2$ and the product $f_1 f_2$ also lie in the Schwartz class. If $f \in \mathcal{S}(\mathbf{R}^n)$, then each translate $f(x - a)$ lies in $\mathcal{S}(\mathbf{R}^n)$ too, and derivatives of f of all orders are also elements of $\mathcal{S}(\mathbf{R}^n)$. Furthermore, the product of f with a polynomial, or a function of the form $\exp(i b \cdot x)$, $b \in \mathbf{R}^n$, lie in $\mathcal{S}(\mathbf{R}^n)$ too.

Let $f(x)$ be a function in the Schwartz class on $\mathbf{R}^n$. Because f is rapidly decreasing, it follows that f is integrable on $\mathbf{R}^n$. Thus we can define the Fourier transform of f as usual by

$$(3) \quad \hat{f}(\xi) = \int_{\mathbf{R}^n} f(x) \exp(-2\pi i \xi \cdot x) dx.$$

Just as for any integrable function on $\mathbf{R}^n$, or any finite measure, the Fourier transform is a bounded continuous function on $\mathbf{R}^n$, and even uniformly continuous. For functions in the Schwartz class we have a much stronger result, namely that the Fourier transform also lies in the Schwartz class.

Indeed, $\hat{f}$ is a bounded continuous function on $\mathbf{R}^n$, and for each multi-index α we have that $x^\alpha f(x)$ is an integrable function on $\mathbf{R}^n$, and hence its Fourier transform is also a bounded continuous function on $\mathbf{R}^n$. Using this one can show that $\hat{f}$ is smooth, and that derivatives of all orders of $\hat{f}$ are bounded. For each multi-index β we have that $\xi^\beta \hat{f}(\xi)$ is a constant times the Fourier transform of $(\partial^{d(\beta)}/\partial x^\beta)f(x)$, which implies that $\xi^\beta \hat{f}(\xi)$ is the Fourier transform of an integrable function, and is therefore bounded. Similarly one can show that derivatives of all orders of $\hat{f}$ decay rapidly, so that $\hat{f} \in \mathcal{S}(\mathbf{R}^n)$.

If ϕ is an integrable function on $\mathbf{R}^n$, then the inverse Fourier transform of ϕ is defined by

$$(4) \quad \check{\phi}(y) = \int_{\mathbf{R}^n} \phi(\eta) \exp(2\pi i y \cdot \eta) d\eta.$$

As for the Fourier transform, the inverse Fourier transform of an integrable function, or more generally a finite measure, is a bounded uniformly continuous function on $\mathbf{R}^n$. If $\phi \in \mathcal{S}(\mathbf{R}^n)$, then $\check{\phi} \in \mathcal{S}(\mathbf{R}^n)$ too, for essentially the same reasons as for the Fourier transform. In general, if f is a continuous integrable function on $\mathbf{R}^n$ such that $\hat{f}$ is also integrable, then the inverse Fourier transform of the Fourier transform of f is equal to f again. In particular, this applies to the case where f is in the Schwartz class, since the Fourier transform of f is then in the Schwartz class as well.

Similarly, if ϕ is a continuous integrable function on $\mathbf{R}^n$ such that $\check{\phi}$ is integrable, then the Fourier transform of $\check{\phi}$ is defined. Under these conditions the Fourier transform of the inverse Fourier transform of ϕ is equal to ϕ , and in particular this holds when $\phi \in \mathcal{S}(\mathbf{R}^n)$.

To summarize a bit, the Fourier transform and inverse Fourier transforms define linear mappings from the Schwartz class into itself which are indeed inverses of each other. It follows that the Fourier transform and inverse Fourier transform are one-to-one linear mappings of the Schwartz class onto itself. Thus every element of the Schwartz class arises as the Fourier transform of an element of the Schwartz class.

If f_1, f_2 are two elements of the Schwartz class on $\mathbf{R}^n$, then the convolu-

tion of f_1 and f_2 is defined by

$$(5) \quad (f_1 * f_2)(x) = \int_{\mathbf{R}^n} f_1(y) f_2(x - y) dy.$$

The Fourier transform of $f_1 * f_2$ is equal to the product of the Fourier transforms of f_1, f_2 . Because f_1, f_2 are elements of the Schwartz class, so are their Fourier transforms. The product of the Fourier transforms of f_1, f_2 therefore lies in the Schwartz class, and one may conclude that $f_1 * f_2$ is an element of the Schwartz class too.

For each function f in the Schwartz class on $\mathbf{R}^n$ and each pair α, β of multi-indices, define $\|f\|_{\alpha, \beta}$ to be the supremum of the absolute value of (1) over all $x \in \mathbf{R}^n$. This defines a seminorm on f , which means that if f_1, f_2 are elements of the Schwartz class, then $\|f_1 + f_2\|_{\alpha, \beta}$ is less than or equal to the sum of $\|f_1\|_{\alpha, \beta}$ and $\|f_2\|_{\alpha, \beta}$, and that if f is an element of the Schwartz class and c is a complex number, then $\|cf\|_{\alpha, \beta}$ is equal to the product of the absolute value of c and $\|f\|_{\alpha, \beta}$. Using this family of seminorms on the Schwartz class we get a topology, which is generated by finite intersections of balls defined with respect to the seminorms.

It is perhaps more convenient to think of this topology in terms of sequences. Namely, if $\{f_j\}_{j=1}^{\infty}$ is a sequence of functions in the Schwartz class and f is another function in the Schwartz class, then $\{f_j\}_{j=1}^{\infty}$ converges to f in the topology determined by the seminorms if and only if $\|f_j - f\|_{\alpha, \beta} \rightarrow 0$ as $j \rightarrow \infty$ for all multi-indices α, β . We can say that a sequence $\{f_j\}_{j=1}^{\infty}$ of functions in the Schwartz class is a Cauchy sequence if $\|f_j - f_l\|_{\alpha, \beta} \rightarrow 0$ as $j, l \rightarrow \infty$ for all multi-indices α, β . One can show that the Schwartz class is complete in the sense that any Cauchy sequence in the Schwartz class converges in the sense just defined to an element of the Schwartz class. The Fourier transform and inverse Fourier transform are continuous linear transformations from the Schwartz class to itself.

The vector space operations of addition and scalar multiplication define continuous mappings from the Schwartz class to itself, as does multiplication of two elements of the Schwartz class. Linear transformations on the Schwartz class defined by differentiating a function in the Schwartz class to some order, or multiplying it by some fixed polynomial, are also continuous. The Schwartz class on $\mathbf{R}^n$ is a basic and natural example of a topological vector space which is a Fréchet space, which means that the topology can be defined by a countable family of seminorms and that the space is complete, while the topology is not determined by any single norm.

By a tempered distribution on $\mathbf{R}^n$ we mean a continuous linear mapping from the Schwartz class on $\mathbf{R}^n$ into the complex numbers. Explicitly, a linear functional λ on $\mathcal{S}(\mathbf{R}^n)$ is continuous if there is a positive real number C and a finite collection of pairs of multi-indices $\alpha_1, \beta_1, \dots, \alpha_l, \beta_l$ such that $|\lambda(f)|$ is less than or equal to C times the sum of $\|f\|_{\alpha_j, \beta_j}$ over $1 \leq j \leq l$ for all functions f in the Schwartz class. A basic class of examples consists of linear functionals λ defined by setting $\lambda(f)$ equal to the integral over $\mathbf{R}^n$ of the product of f and a continuous function h of polynomial growth. If λ is a tempered distribution and α is a multi-index one can define the $\partial^{d(\alpha)}/\partial x^\alpha$ derivative of λ as a tempered distribution by saying that this distribution applied to a function f in the Schwartz class is the same as $(-1)^{d(\alpha)}$ times λ applied to $(\partial^{d(\alpha)}/\partial x^\alpha)f$, which is the same as what one would get by integration by parts if λ is defined by integration of f times a smooth density h . Similarly, one can define the Fourier transform $\hat{\lambda}$ of a tempered distribution by saying that $\hat{\lambda}(f)$ is equal to $\lambda(\hat{f})$ for all f in the Schwartz class.

References

- [1] S. Bochner, *Lectures on Fourier Integrals*, translated by M. Tenenbaum and H. Pollard, Annals of Mathematics Studies **42**, Princeton University Press, 1959.
- [2] W. Rudin, *Functional Analysis*, second edition, McGraw-Hill, 1991.
- [3] L. Schwartz, *Théorie des Distributions*, Tome I et II, Publications de l'Institut de Mathématiques de l'Université de Strasbourg **9** and **10**, Hermann, 1950, 1951.
- [4] E. Stein and G. Weiss, *Introduction to Fourier Analysis on Euclidean Spaces*, Princeton Mathematical Series **32**, Princeton University Press, 1971.